IMAGING COMPLEX STRUCTURES USING BAND LIMITED GREEN'S FUNCTIONS

A DISSERTATION
SUBMITTED TO THE DEPARTMENT OF GEOPHYSICS
AND THE COMMITTEE ON GRADUATE STUDIES
OF STANFORD UNIVERSITY
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

By
Dave Nichols
September 1994

printed as Stanford Exploration Project No. 81
by permission of the author

Preface

All of the figures in this thesis are marked with one of three labels, [R] [NR] or [CR]. These define whether the figure is reproduceable from the information provided on the CD-ROM version of this thesis.

If a figure is marked with an [R] it should be fully reproduceable. All of the input data and source files required to create the figure are provided on the CD. The postscript file containing the figure can be removed and the cakefile rules provided on the CD should rebuild the figure on any of the supported computer architectures.

If a figure is marked with a [CR] it is conditionally reproduceable. All the data and source files are provided on the CD but some particular environment may be necessary to recreate the figure. Many figures in this thesis were created using programs that run on a Thinking Machines Co. CM-5 computer. If the code that is provided will only compile on a CM-5 then the figure will be marked as conditionally reproduceable.

If a figure is marked with a [NR] it is non reproduceable. It cannot be recreated from the data on the CD. The figure may have been created using a commercial software product, or software written by another author that is not available on the CD. It may be a hand drawn figure or a figure for which the commands to recreate it are not available. In all of these cases, if you remove the postscript file of the figure there is no way to recreate it.

All of the reproduceable and conditionally reproduceable figures have a button in the caption. Clicking with the center mouse button will bring up a panel that lets you interact with the figure. You can display the figure in a separate window or destroy and rebuild the figure. If the figure is conditionally reproduceable you can view the warning message that tells you the conditions under which the figure can be rebuilt.

// Abstract

One of the most efficient algorithms for accurate imaging of seismic data is prestack Kirchhoff migration. It has been shown to fail in areas of complex structure, when it is implemented using Green's functions based on first arrival traveltimes. I present a new method for calculating traveltimes that estimates the traveltime of the maximum energy arrival, rather than the first arrival. The estimate is calculated in the seismic frequency band rather than the usual high frequency approximation.

Instead of solving the eikonal equation for the traveltime, I solve the Helmholtz equation to estimate the Green's function for a few frequencies. The Helmholtz equation is solved using a paraxial approximation in a polar coordinate frame. Once a few frequencies have been calculated I perform a parametric fit to the wavefield to estimate a traveltime, amplitude, and phase. The problems associated with aliasing, caused by the coarse sampling in frequency, are avoided by simultaneously extrapolating the wavefield and the traveltime estimates outwards from the source point in the polar coordinate frame.

The images created by using these parametric Green's functions are superior to those created using first arrival traveltimes, or those created by using maximum amplitude traveltimes calculated by paraxial ray tracing. They are comparable in quality to the images created by a full-wavefield downward-continuation scheme.

Acknowledgments

I thank the students, faculty, and sponsors of the Stanford Exploration Project for creating a wonderful environement in which to carry out geophysical research.

I have had many informative discussions over the years with many of my co-workers. Among the most influential were Biondo Biondi, John Etgen, Joe Dellinger, David Lumley, Thorbjorn Rekdal, Matthias Schwab and François Audebert. I would especially like to acknowledge the support of my long time office mates, Steve Cole and Martin Karrenbach. Many problems over the years have been solved with their help and a little shouting across the room.

My advisor, Jon Claerbout, and the other faculty present during my time at SEP, Francis Muir, Fabio Rocca, and Stew Levin, have all helped hone my skills, and keep me going in approximately the correct direction.

I am very grateful to the Institut Français du Pétrole for creating the Marmousi dataset and making it available for comparative research.

Finally I wish to thank my wife and my parents who have always pretended that they believed me when I said I was going to graduate "in a few months". It has really happened at last!

Table of Contents

List of Tables

List of Figures